

## COX-2 INHIBITORS

---

### Products Affected

- Celebrex

<b>Details</b>	
<b>Criteria</b>	<p>Step 1: First line therapy should be a trial and failure of 2 formulary non-selective NSAIDs. NSAID dose must be therapeutic and must be tried for an adequate length of time for anti-inflammatory effects to occur (generally 2-3 weeks).</p> <p>Step 2: Once two of these agents have been tried, patients can receive therapy with Celebrex.</p>

---

# HYPOGLYCEMICS

---

## Products Affected

- Acarbose
- Actoplus Met
- Actoplus Met Xr
- Actos
- Avandamet
- Avandaryl
- Avandia
- Duetact
- Janumet
- Januvia
- Kombiglyze Xr
- Nateglinide
- Onglyza
- Prandin
- Starlix

Details	
<b>Criteria</b>	<p>Step 1: First line therapy should be a formulary sulfonylurea, metformin, or formulary insulin (if appropriate).</p> <p>Step 2: Once one of these agents has been tried, patients can receive therapy with other formulary hypoglycemics such as Actoplus Met, Actoplus Met XR, Actos, Avandamet, Avandaryl, Avandia, Duetact, Acarbose, Januvia, Janumet, Onglyza, Prandin, nateglinide, Kombiglyze XR, and Starlix.</p>

# NON-SEDATING ANTIHISTAMINES

---

## Products Affected

- Allegra
- Clarinex
- Clarinex Reditabs
- Clarinex-d 12 Hour
- Clarinex-d 24 Hour

Details	
<b>Criteria</b>	Step 1: First line therapy should be loratadine OTC (with or without pseudoephedrine) AND cetirizine OTC (with or without pseudoephedrine). Step 2: Second line therapy should be Allegra Suspension. Step 3: Once Allegra Suspension has been tried, patients can receive therapy with other formulary non-sedating antihistamines such as Clarinex.

# OPHTHALMIC ANTIHISTAMINES

---

## Products Affected

- Azelastine Hcl OPHTHALMIC SOLN 0.05%
- Elestat
- Epinastine Hcl
- Optivar
- Pataday
- Patanol

Details	
<b>Criteria</b>	Step 1: First line therapy should be Zaditor OTC, ketotifen OTC or Alaway OTC. Step 2: Second line therapy should be Patanol or Pataday. Step 3: Once Patanol or Pataday has been tried, patients can receive therapy with other formulary prescription ophthalmic antihistamines such as Elestat, azelastine, epinastine, and Optivar.

# PROTON PUMP INHIBITORS

---

## Products Affected

- Aciphex
- Dexilant
- Lansoprazole
- Lansoprazole Odt
- Nexium

Details	
<b>Criteria</b>	Step 1: First line therapy should be Omeprazole or pantoprazole. Step 2: Second line should be lansoprazole or Dexilant. Step 3: Once omeprazole or pantoprazole and lansoprazole or Dexilant has been tried, patients can receive therapy with Nexium or Aciphex.

# INDEX

## **A**

Acarbose.....	2
Aciphex .....	5
Actoplus Met.....	2
Actoplus Met Xr.....	2
Actos.....	2
Allegra.....	3
Avandamet .....	2
Avandaryl.....	2
Avandia .....	2
Azelastine Hcl .....	4

## **C**

Celebrex .....	1
Clarinx .....	3
Clarinx Reditabs .....	3
Clarinx-d 12 Hour.....	3
Clarinx-d 24 Hour.....	3
<b>COX-2 INHIBITORS</b> .....	1

## **D**

Dexilant.....	5
Duetact .....	2

## **E**

Elestat.....	4
Epinastine Hcl .....	4

## **H**

<b>HYPOGLYCEMICS</b> .....	2
----------------------------	---

## **J**

Janumet.....	2
Januvia .....	2

## **K**

Kombiglyze Xr.....	2
--------------------	---

## **L**

Lansoprazole.....	5
Lansoprazole Odt.....	5

## **N**

Nateglinide.....	2
Nexium.....	5

<b>NON-SEDATING ANTIHISTAMINES</b> .....	3
--	---

## **O**

Onglyza.....	2
<b>OPHTHALMIC ANTIHISTAMINES</b> .....	4
Optivar .....	4

## **P**

Pataday.....	4
Patanol.....	4
Prandin .....	2

<b>PROTON PUMP INHIBITORS</b> .....	5
-------------------------------------	---

## **S**

Starlix.....	2
--------------	---